

Supplementary Table 2. Differences in Suicidal Ideation according to General Characteristics

(N = 300)

Variables	Categories	M	SD	t or F	p-value	Scheffé or Games-Howell
Age (yr)	19~29 <sup>a</sup>	23.34	24.14	7.17	.001 <sup>†</sup>	a > c
	30~39 <sup>b</sup>	18.00	23.19			
	40~49 <sup>c</sup>	12.44	16.66			
Marital status	Single	21.37	22.37	2.95	.003 <sup>†</sup>	
	Married	13.99	20.89			
Religion	No	20.19	23.42	2.21	.028 <sup>†</sup>	
	Yes	14.72	19.37			
Education level	High school	22.28	25.76	1.60	.203	
	College	17.12	20.01			
	≥ Graduate school	15.12	25.55			
Chronic disease	No	17.37	21.12	- 1.12	.263	
	Yes	21.55	26.91			
Occupation	No	17.52	25.72	1.31	.270	
	Yes	17.08	20.39			
	Student	23.84	19.35			
Monthly income (10,000 won)	< 100 <sup>a</sup>	21.99	22.04	3.48	.006 <sup>†</sup>	a, b > d
	100~200 <sup>b</sup>	28.76	30.10			
	200~300 <sup>c</sup>	15.82	19.49			
	300~400 <sup>d</sup>	10.46	14.11			
	400~500 <sup>e</sup>	12.22	14.96			
	≥ 500 <sup>f</sup>	15.43	23.40			
Drinking	Drinking <sup>a</sup>	20.42	22.60	3.05	.049	a, c > b
	Drinking in the past <sup>b</sup>	14.15	20.71			
	Never drink <sup>c</sup>	21.21	21.87			
Smoking	Smoking <sup>a</sup>	44.46	31.44	10.89	< .001 <sup>†</sup>	a > b, c
	Smoking in the past <sup>b</sup>	13.48	14.29			
	Never smoke <sup>c</sup>	15.60	19.53			

<sup>†</sup>Welch test (heteroscedasticity test).