

Supplementary Table 3. Differences in Depression and Suicidal Ideation according to COVID-19 Related Characteristics (N = 300)

Variables		M	SD	t	p-value
Depression					
Experience of COVID-19 test	No	6.52	5.26	0.31	.761
	Yes	6.75	5.49		
Have someone close to you been infected by COVID-19?	No	6.50	5.33	0.35	.725
	Yes	6.74	5.22		
Being diagnosed with depression before COVID-19	No	6.11	4.95	3.99	< .001 [†]
	Yes	11.31	6.47		
Taking psychiatric medications prior to COVID-19	No	6.16	5.04	4.98	< .001
	Yes	11.90	5.84		
Have attempted suicide before COVID-19	No	6.39	5.12	3.69	< .001
	Yes	13.71	7.70		
Suicidal ideation					
Experience of COVID-19 test	No	17.83	21.10	0.15	0.878
	Yes	18.32	25.37		
Have someone close to you been infected by COVID-19?	No	16.74	20.98	1.54	0.126
	Yes	21.12	24.30		
Being diagnosed with depression before COVID-19	No	15.96	19.75	5.24	< .001
	Yes	38.62	32.01		
Taking psychiatric medications prior to COVID-19	No	16.15	19.39	3.12	.005 [†]
	Yes	41.52	36.93		
Have attempted suicide before COVID-19	No	16.48	19.07	3.66	.010 [†]
	Yes	78.43	44.65		

[†]Welch test (heteroscedasticity test).