

Supplementary Table 5. Factors Associated with Suicidal Ideation

(N = 300)

Variables	B	SE	$\beta$	t	p-value
Constant	6.77	7.45		0.91	.364
Age (yr)					
30~39	- 1.58	2.37	- 0.03	- 0.67	.505
40~49	- 5.01	2.82	- 0.11	- 1.78	.076
Marital status (married)	0.97	2.25	0.02	0.43	.666
Religion (yes)	- 4.10	1.78	- 0.09	- 2.30	.022
Monthly income (10,000 won)					
100~200	3.25	3.07	0.05	1.06	.290
200~300	- 2.35	2.52	- 0.05	- 0.94	.351
300~400	- 0.45	3.65	- 0.01	- 0.12	.903
400~500	- 2.17	3.52	- 0.03	- 0.62	.537
≥ 500	- 0.90	3.23	- 0.01	- 0.28	.781
Drinking					
Drinking	- 5.87	4.21	- 0.13	- 1.39	.164
Drinking in the past	- 7.25	4.27	- 0.16	- 1.70	.090
Smoking					
Smoking	17.04	3.12	0.22	5.47	< .001
Smoking in the past	- 0.27	3.10	0.00	- 0.09	.930
Being diagnosed with depression before COVID-19 (yes)	3.74	3.89	0.05	0.96	.337
Taking psychiatric medications prior to COVID-19 (yes)	3.35	4.30	0.04	0.78	.437
Having attempted suicide before COVID-19 (yes)	39.68	5.82	0.27	6.82	< .001
Social distancing	0.02	0.15	0.01	0.14	.890
Fear of COVID-19	0.76	0.76	0.04	1.00	.318
Depression	2.05	0.19	0.49	11.09	< .001
F (p)			21.84 (< .001)		
Adjusted R <sup>2</sup>			.57		
d (d <sub>u</sub> )			2.03 (1.95)		

d (d<sub>u</sub>) = Durbin-Watson's auto-correlation coefficient (upper critical limit).

Reference groups: age (19~29 yr), marital status (single), monthly income (10,000 won, < 100), drinking (never drink), smoking (never smoke), being diagnosed with depression before COVID-19 (no), taking psychiatric medications prior to COVID-19 (no), having attempted suicide before COVID-19 (no).